

**RONICOX<sup>®</sup>**  
**Powder for oral administration**

**Description**

Light brown powder.

**Composition**

100 g of the drug contains:

*active ingredient*: robenidine hydrochloride – 6.6 g;

*excipient* - semolina.

**Pharmacological properties**

*ATCvet*: **QP51**. Antiprotozoals. **QP51AX13**. Robenidine.

*Robenidine hydrochloride* is eimeriostatic, which is effective against eimeriae of the following species: *Eimeria necatrix*, *E. tenella*, *E. acervulina*, *E. maxima*, *E. adenoeides*, *E. gallopavonis*, *E. meleagriditis* at the stage of schizonts of the first and second generations.

It quickly passes the digestive tract of poultry and has a low absorption level (20-40%). It is primarily eliminated with feces. The elimination volume makes 60-65% of the administered dose.

**Administration**

Treatment and prevention of *Eimeria* lesions in broiler chickens and turkeys:

- **broiler chickens** - *Eimeria tenella*, *E. acervulina*, *E. mivati*, *E. brunetti*, *E. maxima*, *E. necatrix*;

- **turkeys (up to 16 weeks of age)** - *Eimeria adenoeides*, *E. meleagriditis* and *E. gallopavonis*.

**Dosage**

Ronicox<sup>®</sup> is fed to broiler chickens and turkeys with feed in a dose of 450-500 g per 1 t of the feed, which corresponds to 30-33 g of robenidine hydrochloride per 1 t of the feed.

The drug is administered to turkeys from the first days of life up to 16 weeks, to broiler chickens – from the first days of life throughout the growing period.

**Contraindications**

Do not administer to laying hens.

**Precautions**

Animal slaughter for meat is possible in 5 days following the last administration. Meat obtained before the mentioned term shall be utilized or fed to non-productive animals depending on the statement of veterinary physician.

When working with the drug it is recommended that protective clothes, respirator, gloves and tightly fitting goggles be used. In case of contact with eyes wash them with plenty of water for 15 minutes.

**Packaging**

Bags of heat sealable film materials or double PET-bags of 1 000 g; multilayer paper bags of 25 kg.

**Storage**

Store in a dry, dark place out of the reach of children at 15-30°C.

**Shelf life**

2 years.