

UNISAN[®] Ointment

Description

White ointment with specific odor.

Composition

1 g of the drug contains:

active ingredients: chlorhexidine bigluconate – 10 mg; zinc oxide – 50 mg;

excipients - dimethyl sulfoxide, PEG-1500, distilled water.

Pharmacological properties

ATCvet: *QD08AC52*. Chlorhexidine, combinations.

The drug has a pronounced antiseptic effect. The components of ointment are effective against vegetative forms of gram-positive (*Staphylococcus spp.*, *Streptococcus spp.*, *Mycobacterium spp.*, *Corynebacterium xerosis*, *Micrococcus spp.*, *Sarcina lutea*, *Clostridium spp.*, *Bacillus subtilis*) and gram-negative bacteria (*Escherichia coli*, *Salmonella spp.*, *Shigella spp.*, *Klebsiella spp.*, *Enterobacter spp.*, *Serratia marcescens*, *Providencia spp.*, *Proteus spp.*, *Pseudomonas spp.*, *Neisseria spp.*) as well as yeasts (*Candida spp.*), dermatophytes (*Dermatophytes*, *Microsporium*, *Trichophyton*, *Trichoderma*), lipophilic viruses, chlamydiae. Moreover, the drug makes anti-inflammatory, astringent, local analgesic effect.

Administration

Apply externally to treat skin diseases caused by bacterial and fungal microflora, that is: skin staphylococcus, pyoderma, impetigo eczema, skin candidosis, trichophytosis, microsporia, seborrhea, external otitis, septic wounds, superficial abscesses and burns in cattle, sheep, goats, horses, swine, rabbits, poultry, dogs and cats.

Dosage

The ointment is applied with a thin layer on the injured skin area once or twice daily until complete recovery. Treatment period is conditioned by the type of pathological process. In any case it is necessary to apply ointment for at least 2-3 days after disappearance of clinical features of disease. In case of fungal diseases continue treatment for 7-10 days after disappearance of clinical symptoms of disease. To treat septic wounds and abscesses, permeate gauzes with the ointment and apply the gauzes to wounds or abscess cavities. Apply dressings every day until complete recovery from septic and necrotic masses.

Contraindications

Do not apply in case of individual hypersensitivity to chlorhexidine. Avoid contact with mucous membranes. Do not prescribe in combination with iodine drugs.

Precautions

Utilize unused product in accordance with requirements of national legislation.

Packaging

Polymer jars, vials and aluminum tubes of 15 g and 200 g.

Storage

Store in a dark place out of the reach of children at 0-25°C.

Shelf life

2 years.